



History:

Received: October 12, 2015
Accepted: November 1, 2015
First Published: January 1, 2016
Collection year: 2015
Confirmation of publication: Published

Identifiers and Pagination:

Year: 2016
Volume / Issue: 8/1
First Page: 7
Last Page: 10
Publisher Id: JAppPharm-8-1
DOI:<http://dx.doi.org/10.21065/19204159>

Corresponding author:

Shaukat Khanum Memorial Cancer Hospital & Research Center 7 A Block R3 Johar Town Lahore, Pakistan.
Tel.: +92-332-6070971.
E. adeels@live.com

Citation:

Adeel Siddiqui, Zuhaib Jaffer Malik, Sidrah Andleeb. Vancomycin associated eosinophilia & red man syndrome in a tertiary care hospital. J App Pharm (2016) 8:1. 7-10

Original Research Article

VANCOMYCIN ASSOCIATED EOSINOPHILIA & RED MAN SYNDROME IN A TERTIARY CARE HOSPITAL

Adeel Siddiqui^{1,2}, Zuhaib Jaffer Malik³, Sidrah Andleeb³

1. Department of Pharmacy, Shaukat Khanum Memorial Cancer Hospital & Research Center, Lahore, Pakistan
2. Department of Pharmacy, Quaid-i-Azam University, Islamabad, Pakistan
3. Department of Pharmacy, Shaukat Khanum Memorial Cancer Hospital & Research Center, Lahore, Pakistan

Abstract

Vancomycin associated Red Man Syndrome (RMS) is a common infusion related adverse effect. A correlation has been proposed between vancomycin induced eosinophilia and occurrence of RMS. In such cases, it is identified that activated T-cells release cytokines, which tends to contribute to hypersensitivity reaction. A cross-sectional, retrospective study was conducted to evaluate the possible association of vancomycin induced eosinophilia with the occurrence of RMS at a local tertiary care cancer hospital. In this study the role of eosinophilia before the occurrence of RMS was focused. A twenty one month data was extracted from Jan, 2014 to Sep, 2015 and 8 patients were found to be reported for vancomycin induced RMS. Only one patient (12.5%) fulfilled the criteria for the drug rash eosinophilia and systemic symptoms (DRESS) among the 8 patients. This particular patient had eosinophilia preceding RMS. In the study sample, 2(25%) patients showed a rise in their eosinophil count but within the normal range. A correlation of eosinophilia and RMS could not be established in our study.

Keywords: red man syndrome, vancomycin, drug rash eosinophilia and systemic symptoms, eosinophilia.

Introduction

Vancomycin is a significant treatment option for Methicillin-resistant *Staphylococcus aureus* (MRSA) infection, and a first line therapy for Methicillin Resistant *Streptococcus pneumoniae* and *Corynebacterium jeikeium* infection [1-3]. For the treatment of methicillin-susceptible (MSSA) infections it is less preferred relative to beta-lactams [1-2].

In spite of an effective Antimicrobial Stewardship Program (ASP) carried out in different hospitals, the use of Vancomycin was increasing due to increased prevalence of MRSA cases. This may have happened due to irrational prescribing of antibiotics in third world countries leading to multi-drug resistance [4]. Thus, in patient receiving vancomycin it's important to establish the complications in antibiotic therapy from occurrence of DRESS.

Use of Vancomycin has led to two types of hypersensitivity reaction, one is the anaphylaxis and other is RMS. The latter is associated with face, neck and abdomen involvement, with prevalence of pruritus and rash. Patients tend to experience a feeling of burning, leading to discomfort. Symptoms also include headache, hyperthermia, itching and paresthesia, due to which patient becomes agitated [5].

Incidence of allergic reaction may also serve for clinical diagnosis of DRESS following drug use. The criteria for DRESS include hyperthermia, rash, eosinophilia, and existence of multi-organ association

Reviewing editor:

Taha Nazir, Ph.D. Scientific Executive
ICDTD Inc. Saskatoon Saskatchewan
Canada.: E. taha@icdti.ca

Funding:

The authors received no direct
funding for this research.

Competing Interests:

The authors declare no competing
interests

Additional information is available at
the end of the article.

[6]. Other hematological changes may include leukocytosis or atypical lymphocytes. The usual organ involvement tends to appear within 2 weeks of skin erythema [7-8]. In RMS eosinophilia generally occurs, however, in case of vancomycin induced RMS it is important to distinguish the route cause. Generally, RMS is correlated to the higher infusion rates. Thus, the claim that eosinophilia is the cause for RMS may be questionable.

A study was conducted at Shaukat Khanum Memorial Cancer Hospital & Research Center, Lahore (SKMCH&RC), Pakistan, a local tertiary care hospital, with the aim to assess the significance of eosinophilia in patients experiencing RMS with vancomycin use.

MATERIALS AND METHODS

A retrospective cross-sectional study was conducted to evaluate the adverse drug reactions in terms of Vancomycin induced RMS, reported from Jan, 2014 to Sep, 2015. A total of 2,666 patient data gathered, who received vancomycin during the sample period.

Results

In the study sample (n= 2666) 8 patients were reported to suffer from vancomycin induced RMS. Among these patients, 2(25%) patients had their eosinophil count slightly deviated but it was within normal range and showed rise in eosinophil count. Only 1(12.5%) patient had prevalence of eosinophilia along with RMS, fulfilling the criteria of DRESS. The data collected of patients who had RMS post-vancomycin, is depicted in Figure 1.




Figure 1: Data for RMS from Jan, 2014 to Sep, 2015

Discussion

In this study correlation of eosinophilia with occurrence of RMS in patients receiving vancomycin was evaluated. Eosinophilia may serve as a marker to the definitive occurrence of RMS and may facilitate the physician as evidence to discontinue the specific drug. In terms of the mechanism, RMS is a result of type IV hypersensitivity reaction due to cytokines released for the activated T-cells [9].

DRESS was earlier connected with the use of anticonvulsants [10-11], but now an increasing trend has been noticed for it in patients receiving antibiotics such as, β -lactams, vancomycin, sulfa-drugs, tetracycline and flouroquinolone. In our study the focus was on vancomycin.

Vancomycin use is now increasing in hospitals as a result of growing number of multidrug resistant bacteria. In case of incidence of RMS, linezolid substitution is considered as one of the options. If the incidence is independent of eosinophilia then the most probable cause can be high rate of infusion, in which case the drug may be continued at a slower rate. The usual treatment for RMS is antihistamines and intravenous or topical corticosteroids.

A similar research conducted by KG Blumenthal et al showed a 4 years data collection for vancomycin induced DRESS, where 5 patients fulfilled the criteria. Comparative to this, our study had 1 patient, who had eosinophilia and fulfilled the criteria for DRESS [12].

neutropenia started with Piperacillin/Tazobactam 1900mg q6h as an empiric therapy. Later the patient developed fever and blood culture exhibited gram-positive *Micrococcus species* against which vancomycin was started. The patients' hematological test showed raised eosinophil count which consistently escalated as the therapy proceeded. On day 8 of vancomycin, he started to develop rashes with typical signs and symptoms of RMS, therefore, Vancomycin was stopped at once. Hydrocortisone and Pheniramine were administered for symptomatic management. Condition settled soon after vancomycin was discontinued and it was replaced with teicoplanin.

Two pediatric cases were discussed by Bauters T, et al [13] with cancer malignancy, receiving vancomycin. Both patients were infused drug over 60 minutes. Interestingly, none were on opioids, but they discussed the role of concomitant use of opioids in the occurrence of RMS [14]. In our case, Patient was receiving tramadol for pain management.

Conclusion

It was concluded that vancomycin may not influence eosinophil count. Also, raised eosinophil count is unlikely to contribute to RMS in patients treated with vancomycin. Further studies are recommended to establish the incidence of RMS secondary to eosinophilia in vancomycin treated patients.

REFERENCES

1. Kollef MH. (2007). Limitations of vancomycin in the management of resistant staphylococcal infections. *Clin Infect Dis* 45: S191-5
2. Stryjewski ME, Szczech LA, Benjamin DK Jr (2007). Use of Vancomycin or first-generation cephalosporins for the treatment of hemodialysis-dependent patients with methicillin-susceptible *Staphylococcus aureus* bacteremia. *Clin Infect Dis*. 44(2):190
3. Eggimann P, Pittet D (2001). Infection control in the ICU. *Chest* 120:2059-2093
4. Michael Z. David, Robert S. Daum (2010). Community –Associated Methicillin-Resistant *Staphylococcus aureus*: Epidemiology and Clinical Consequences of an Emerging Epidemic. *Clin Microbiol* 23(3): 616-687
5. Davis RL, Smith AL, Koup JR (1986). The 'red man's syndrome' and slow infusion of vancomycin. *Ann Intern Med*. 104:285-286
6. Kennbeck GA (2000). Anticonvulsant hypersensitivity syndrome. *J Am Board Fam Pract* 13:364-370

7. Ganceva M, Ganceva T, Lazarova R (2008). Carbamazepine-induced drug reaction with eosinophilia and systemic symptoms (DRESS) syndrome: Report of four cases and brief review. *Int J Dermatol* 47:853-860
8. Ben m'rad M, Leclerc-Mercier S, Blanche P (2009). Drug induced hypersensitivity syndrome. *Medicine* 88:131-140
9. Gell PGH, Coombs RRA (1963). The classification of allergic reactions underlying disease. *Clinical Aspects of Immunology*, 1st ed. Oxford, England: Blackwell, 317-337
10. Cacoub P, Musette P, Descamps V, Meyer O, Speirs C, Finzi L (2011). The DRESS syndrome: a literature review. *Am J Med* 124:588-97
11. Jeung J, Lee JY, Oh MJ, Choi DC, Lee BJ (2010). Comparison of the causes and clinical features of drug rash with eosinophilia and systemic symptoms and Steven Johnson's syndrome. *Allergy Asthma Immunol Res* 2:123-6
12. Blumenthal KG, Patil SU, Long AA (2012). The importance of vancomycin in drug rash with eosinophilia and systemic symptoms (DRESS) syndrome. *Allergy Asthma Proc* 33:165-171
13. Bauters T, Claus B, Schelstraete P, Robays H, Benoit Y, Dhooge C (2012). Vancomycin-induced red man syndrome in pediatric oncology: still an issue? *34:13-6*
14. Levy JH, Marty AT. Vancomycin and adverse drug reactions. *Crit Care Med*. 1993;21(8):1107-8.

The single reported patient diagnosed T-cell Non-Hodgkin lymphoma, admitted with febrile



© 2016 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits